

The First and Second Laws of Gravity

Maulik Parikh

Department of Physics
Arizona State University

M. P. and A. Svesko, “The First Law of Gravity,” to appear.

M. P. and A. Svesko, “Thermodynamic Origin of the Null Energy Condition,” arXiv: 1511.06460; PLB.

M. P. and J.-P. van der Schaar, “Derivation of the Null Energy Condition,” arXiv: 1406.5163; Phys. Rev. D.

Science or Science Fiction?

Gödel universe

Gott time machine

Morris-Thorne wormhole


Bouncing cosmologies

Big Rip phantom cosmology

Negative mass black holes

Super-extremal black holes

Einstein on the Gödel Universe



“... cosmological solutions of the gravitation equations have been found by Mr Gödel. It will be interesting to weigh whether these are not to be excluded on physical grounds.”

“Physical Grounds” or Ad Hoc Criteria?

No closed causal curves

Stable causality

Global hyperbolicity

Cosmic censorship/strong asymptotic predictability

Geodesic completeness

Generalized second law of thermodynamics

Energy conditions

The Null Energy Condition

The most basic of the energy conditions is the null energy condition (NEC):

$$T_{\mu\nu}k^{\mu}k^{\nu} \geq 0$$

where k is any light-like vector.

The weak and strong energy conditions both imply the null energy condition.

Importance of Energy Conditions

Singularity theorems.

Positive energy theorems.

Black hole no-hair theorem.

Laws of black hole mechanics.

Exclusion of traversable wormholes, construction of time machines, creation of a universe in the laboratory, and cosmological bounces.

$$\dot{H} = -(\rho + p) \leq 0$$

Energy Conditions from Matter?

The energy conditions are conventionally viewed as conditions on the energy-momentum tensor for *matter*.

Matter

We have a fantastic framework for understanding matter: quantum field theory.

Violation of the Null Energy Condition in QFT

Consider a well-behaved NEC-obeying theory of matter, theory A.

Let theory B have the same action as theory A but with an overall minus sign.

Theory B then violates the NEC.

But theory B is otherwise just as well-behaved as theory A!

Hence the NEC does not follow from QFT.

How the Energy Conditions are Used in Practice

In gravitational theorems, the energy conditions are always used in conjunction with the equations of motion.

$$T_{\mu\nu}k^\mu k^\nu \geq 0 \Rightarrow (R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu})k^\mu k^\nu \geq 0 \Rightarrow R_{\mu\nu}k^\mu k^\nu \geq 0$$

Why not just start with a condition on the Ricci tensor?

Perhaps the real null “energy condition” is

$$R_{\mu\nu}k^\mu k^\nu \geq 0$$

We will derive this condition.

Viewpoint in this Talk

We will take

$$R_{\mu\nu}k^{\mu}k^{\nu} \geq 0$$

as a fundamental property of gravity. We will continue to refer to this as the “null energy condition”. But note that this is now a condition on geometry.

Classical Gravity

Property 1:

$$G_{\mu\nu} = T_{\mu\nu}$$

Property 2:

$$R_{\mu\nu} k^\mu k^\nu \geq 0$$

Local Holographic Thermodynamics

Premise of Talk

Associate gravitational entropy *locally* to patches of certain null surfaces.

Assume that this entropy arises from the coarse-graining of some dual microscopic statistical-mechanical system.

Emergent Gravity

Jacobson (1995) showed that Einstein's equations followed from the Clausius relation $dQ = TdS$ when applied to "local Rindler horizons".

Our prescription will be somewhat different. We will attribute thermodynamic properties to infinitesimal (patches of) expanding future light cones.

First Law

Generalized Einstein Equations

Consider an arbitrary relativistic diffeomorphism-invariant theory of gravity:

$$I = \frac{1}{16\pi} \int d^D x \sqrt{-g} L(g_{ab}, R_{abcd}) + I_{\text{matter}}$$

Define

$$P^{abcd} = \frac{\partial L}{\partial R_{abcd}}$$

Then the equation of motion of classical gravity is

$$P_a{}^{cde} R_{bcde} - 2\nabla^c \nabla^d P_{acdb} - \frac{1}{2} L g_{ab} = 8\pi T_{ab}$$

For example, for Einstein gravity, $L = R$, and $P^{abcd} = \frac{1}{2}(g^{ac}g^{bd} - g^{ad}g^{bc})$

and this reduces to Einstein's equations.

Goal: Gravitational Equations from the First Law

Our goal is to derive the generalized Einstein equations

$$P_a^{cde} R_{bcde} - 2\nabla^c \nabla^d P_{acdb} - \frac{1}{2} L g_{ab} = 8\pi T_{ab}$$

from the first law of thermodynamics, $dM = TdS$, applied locally.

We need to define M , T , and S .

Local Killing Vectors

Consider an arbitrary point, P , in an arbitrary spacetime.

In the vicinity of P , spacetime is approximately flat.


Hence there are local Lorentz symmetries, generated by approximate Killing vectors.

M

For a Killing vector ξ_a

$$\nabla_a (T^{ab} \xi_b) = 0$$

Integrating over a compact region of spacetime, we have



$$M_i \equiv \int_{S_i} d^{D-1}x \sqrt{h} u_a T^{ab} \xi_b$$

$$\Delta M = M_2 - M_1 = \int_{\Sigma} d\Sigma_a T^{ab} \xi_b$$

T

In the vicinity of P, spacetime is approximately flat. The local Lorentz symmetries include boosts. Consider a surface generated by *radial* boosts.

$$\xi^a = r\partial_t^a + t\partial_r^a$$

The surface $\xi^2 = -\alpha^2$ is a timelike de Sitter-like hyperboloid. An observer moving along a boost has a constant proper acceleration, $1/\alpha$.

Define

$$T = \frac{1}{2\pi\alpha}$$

S

Wald found an expression for black hole entropy in general theories of gravity:


$$S = \frac{\alpha}{8} \int_S dS_{ab} J^{ab}$$

where $J^{ab} = -2P^{abcd}\nabla_c\xi_d + 4\xi_d(\nabla_c P^{abcd})$

is the “Noether potential” for a timelike Killing vector, ξ_a .

The Wald entropy reduces to $A/4$ for Einstein gravity.

Change in Entropy due to Infalling Matter



$$\Delta S = \Delta S_{\text{tot}} - \Delta S_0$$

Variation of Wald Entropy

Apply the Wald entropy formula to our infinitesimal compact surface:

$$S_{\text{tot}} = \frac{\alpha}{8} \int_S dS_{ab} J^{ab} = -\frac{\alpha}{4} \int_S dS_{ab} (P^{abcd} \nabla_c \xi_d - 2\xi_d \nabla_c P^{abcd})$$

After some manipulation, we find

$$\Delta S_{\text{tot}} = +\frac{\alpha}{4} \int_{\Sigma} d\Sigma_a [P^{abcd} R_{bcd}^e \xi_e - 2\xi_d \nabla_b \nabla_c P^{abcd}] + \Delta S_0$$

First Law of Gravity

Therefore the first law of gravity says

$$\Delta M = T \Delta S = T(\Delta S_{\text{tot}} - \Delta S_0)$$

where $\Delta M = \int_{\Sigma} d\Sigma_a T^{ab} \xi_b$ and $T = \frac{1}{2\pi\alpha}$

and

$$\Delta S = \frac{\alpha}{4} \int_{\Sigma} d\Sigma_a (P^{acde} R_{cde}^b - 2\nabla_c \nabla_d P^{acdb} \xi_b)$$

Putting this all together (and using the Bianchi identity) we find Einstein's equations:

$$P_a^{cde} R_{bcde} - 2\nabla^c \nabla^d P_{acdb} - \frac{1}{2} L g_{ab} = 8\pi T_{ab}$$

Second Law

From the Second Law to the NEC

Quote a result about near-equilibrium thermodynamic systems obeying the second law.

Attribute thermodynamic properties to future light cones.

Show that null congruences corresponding to near-equilibrium thermodynamic systems exist at every point.

Obtain the null energy condition from the second law.

Onsager Theory


Near-equilibrium thermodynamic systems
approaching internal equilibrium via the second
law obey not only

$$\dot{S} \geq 0$$

but also

$$\ddot{S} \leq 0$$

G. Falkovich and A. Fouxon, New J. Phys. 6 (2004) 50.



Entropy and Area

As before, attribute Bekenstein-Hawking entropy to infinitesimal patches of expanding future light cones (perhaps more generally, to all non-contracting infinitesimal patches of future-directed null congruences).

Then

$$\begin{aligned} S &= \frac{A}{4} \\ \dot{S} &= \frac{A}{4} \theta \\ \ddot{S} &= \frac{A}{4} (\dot{\theta} + \theta^2) \end{aligned}$$

For patches corresponding to near-equilibrium systems, the last term is non-positive.

Null Energy Condition from Thermodynamics

The infinitesimal patch obeys the Raychaudhuri equation:

$$\dot{\theta} = -\frac{1}{2}\theta^2 - R_{\mu\nu}k^\mu k^\nu$$

Rewriting that equation we find, for near-equilibrium congruences, that

$$\begin{aligned} R_{\mu\nu}k^\mu k^\nu &= -\dot{\theta} - \frac{1}{2}\theta^2 \\ &= -\left(\dot{\theta} + \theta^2\right) + \frac{1}{2}\theta^2 \\ &= -\frac{1}{S}\ddot{S} + \frac{1}{2S^2}\dot{S}^2 \\ &\geq 0 \end{aligned}$$

This is precisely the geometric form of the NEC.

The First and Second Laws of Gravity

First law:

$$G_{\mu\nu} = T_{\mu\nu}$$

Second law:

$$R_{\mu\nu} k^{\mu} k^{\nu} \geq 0$$

Stringy Origin of the Null Energy Condition

Strings in Curved Spacetime

For a string propagating in curved space, the one-loop effective action is

$$S[X_0^\mu, h_{ab}] = -\frac{1}{4\pi\alpha'} \int d^2\sigma \sqrt{-h} h^{ab} \partial_a X_0^\mu \partial_b X_0^\nu (\eta_{\mu\nu} + C_\epsilon \alpha' R_{\mu\nu}(X_0)) - \frac{1}{4\pi} \int d^2\sigma \sqrt{-h} C_\epsilon \Phi(X_0) R^{(2)}$$

Virasoro constraint in light-cone coordinates:

$$0 = \partial_+ X_0^\mu \partial_+ X_0^\nu (\eta_{\mu\nu} + C_\epsilon \alpha' R_{\mu\nu} + 2C_\epsilon \alpha' \nabla_\mu \nabla_\nu \Phi)$$

At zeroth order in α' we find that

$$\eta_{\mu\nu} \partial_+ X^\mu \partial_+ X^\nu = 0$$

In other words, $k^\mu \equiv \partial_+ X^\mu$ is a *spacetime* null vector field.

But at first order in α' we miraculously discover that

$$k^\mu k^\nu (R_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi) = 0$$

Einstein-Frame Metric

The spacetime metric appearing in the string action is the string-frame metric.

It is related to the usual (“Einstein-frame”) metric by scaling:

$$g_{\mu\nu} = e^{+\frac{4\Phi}{D-2}} g_{\mu\nu}^E$$

Transforming the Virasoro condition to the Einstein-frame metric, we find





$$0 = k^\mu k^\nu (R_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi) = k^\mu k^\nu \left(R_{\mu\nu}^E - \frac{4}{D-2} \nabla_\mu^E \Phi \nabla_\nu^E \Phi \right)$$

That is

$$R_{\mu\nu}^E k^\mu k^\nu = + \frac{4}{D-2} (k^\mu \nabla_\mu^E \Phi)^2 \geq 0$$

This is precisely the desired geometric form of the null energy condition!

Local Holography?

Closed String Worksheet	Spacetime	Holographic Thermodynamics
Weyl invariance 	Einstein's equations	 First law
Diffeomorphism invariance 	Null energy condition	 Second law