

# Obstructions to embeddings into $L_p$ spaces: Property $\alpha$

Gideon Schechtman

Joint work with  
Assaf Naor

BIRS, May 2017

# Bi-Lipschitz embedding

A metric space  $(X, d_X)$  is said to admit a bi-Lipschitz embedding into a metric space  $(Y, d_Y)$  if there exist  $s \in (0, \infty)$ ,  $D \in [1, \infty)$  and a mapping  $f : X \rightarrow Y$

$$\forall x, y \in X, \quad sd_X(x, y) \leq d_Y(f(x), f(y)) \leq Dsd_X(x, y).$$

When this happens we say that  $(X, d_X)$  embeds into  $(Y, d_Y)$  with distortion at most  $D$ . We denote by  $c_Y(X)$  the infimum over such  $D \in [1, \infty]$ . When  $Y = L_p$  we use the shorter notation  $c_{L_p}(X) = c_p(X)$ .

We will be interested in bounding from below the distortion of embedding certain metric spaces into  $L_p$ . I'll concentrate on embedding certain grids in Schatten  $p$ -classes into  $L_p$ .

# Bi-Lipschitz embedding

A metric space  $(X, d_X)$  is said to admit a bi-Lipschitz embedding into a metric space  $(Y, d_Y)$  if there exist  $s \in (0, \infty)$ ,  $D \in [1, \infty)$  and a mapping  $f : X \rightarrow Y$

$$\forall x, y \in X, \quad sd_X(x, y) \leq d_Y(f(x), f(y)) \leq Dsd_X(x, y).$$

When this happens we say that  $(X, d_X)$  embeds into  $(Y, d_Y)$  with distortion at most  $D$ . We denote by  $c_Y(X)$  the infimum over such  $D \in [1, \infty]$ . When  $Y = L_p$  we use the shorter notation  $c_{L_p}(X) = c_p(X)$ .

We will be interested in bounding from below the distortion of embedding certain metric spaces into  $L_p$ . I'll concentrate on embedding certain grids in Schatten  $p$ -classes into  $L_p$ .

# Schatten classes

Given a (finite or infinite, real or complex) matrix  $A$  and  $1 \leq p < \infty$

$$\|A\|_p = (\text{trace}(A^*A)^{p/2})^{1/2} = \left(\sum_{i=1}^{\infty} \lambda_i^p\right)^{1/p}$$

where the  $\lambda_i$ -s are the singular values of  $A$ .

$$\|A\|_{\infty} = \|A : \ell_2 \rightarrow \ell_2\|.$$

$S_p^n$  is the space of all  $n \times n$  matrices equipped with the norm  $\|\cdot\|_p$ .

$e_{ij}$  denotes the matrix with 1 in the  $ij$  place and zero elsewhere. This is a good basis in a certain order but, except if  $p = 2$ , NOT a good unconditional basis.

Here is a simple way to prove it:

# Schatten classes

Given a (finite or infinite, real or complex) matrix  $A$  and  $1 \leq p < \infty$

$$\|A\|_p = (\text{trace}(A^*A)^{p/2})^{1/2} = \left(\sum_{i=1}^{\infty} \lambda_i^p\right)^{1/p}$$

where the  $\lambda_i$ -s are the singular values of  $A$ .

$$\|A\|_{\infty} = \|A : \ell_2 \rightarrow \ell_2\|.$$

$S_p^n$  is the space of all  $n \times n$  matrices equipped with the norm  $\|\cdot\|_p$ .

$e_{ij}$  denotes the matrix with 1 in the  $ij$  place and zero elsewhere. This is a good basis in a certain order but, except if  $p = 2$ , NOT a good unconditional basis.

Here is a simple way to prove it:

# Schatten classes

Given a (finite or infinite, real or complex) matrix  $A$  and  $1 \leq p < \infty$

$$\|A\|_p = (\text{trace}(A^*A)^{p/2})^{1/2} = \left(\sum_{i=1}^{\infty} \lambda_i^p\right)^{1/p}$$

where the  $\lambda_i$ -s are the singular values of  $A$ .

$$\|A\|_{\infty} = \|A : \ell_2 \rightarrow \ell_2\|.$$

$S_p^n$  is the space of all  $n \times n$  matrices equipped with the norm  $\|\cdot\|_p$ .

$e_{ij}$  denotes the matrix with 1 in the  $ij$  place and zero elsewhere. This is a good basis in a certain order but, except if  $p = 2$ , NOT a good unconditional basis.

Here is a simple way to prove it:

Given a (finite or infinite, real or complex) matrix  $A$  and  $1 \leq p < \infty$

$$\|A\|_p = (\text{trace}(A^*A)^{p/2})^{1/2} = \left(\sum_{i=1}^{\infty} \lambda_i^p\right)^{1/p}$$

where the  $\lambda_i$ -s are the singular values of  $A$ .

$$\|A\|_{\infty} = \|A : \ell_2 \rightarrow \ell_2\|.$$

$S_p^n$  is the space of all  $n \times n$  matrices equipped with the norm  $\|\cdot\|_p$ .

$e_{ij}$  denotes the matrix with 1 in the  $ij$  place and zero elsewhere. This is a good basis in a certain order but, except if  $p = 2$ , NOT a good unconditional basis.

Here is a simple way to prove it:

# Schatten classes

Given a (finite or infinite, real or complex) matrix  $A$  and  $1 \leq p < \infty$

$$\|A\|_p = (\text{trace}(A^*A)^{p/2})^{1/2} = \left(\sum_{i=1}^{\infty} \lambda_i^p\right)^{1/p}$$

where the  $\lambda_i$ -s are the singular values of  $A$ .

$$\|A\|_{\infty} = \|A : \ell_2 \rightarrow \ell_2\|.$$

$S_p^n$  is the space of all  $n \times n$  matrices equipped with the norm  $\|\cdot\|_p$ .

$e_{ij}$  denotes the matrix with 1 in the  $ij$  place and zero elsewhere. This is a good basis in a certain order but, except if  $p = 2$ , NOT a good unconditional basis.

Here is a simple way to prove it:



# Schatten classes

For simplicity,  $p = 1$ .

## Claim

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{e}_{ij} \right\|_1 \approx n^{3/2},$$

While

$$\left\| \sum_{i,j=1}^n \mathbf{e}_{ij} \right\|_1 = n.$$

The  $\geq$  side in the first equivalence follows easily from duality between  $S_1^n$  and  $S_\infty^n$  and the not-hard fact that

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{e}_{ij} \right\|_\infty \lesssim n^{1/2}.$$

# Schatten classes

For simplicity,  $p = 1$ .

Claim

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{e}_{ij} \right\|_1 \approx n^{3/2},$$

While

$$\left\| \sum_{i,j=1}^n \mathbf{e}_{ij} \right\|_1 = n.$$

The  $\geq$  side in the first equivalence follows easily from duality between  $S_1^n$  and  $S_\infty^n$  and the not-hard fact that

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{e}_{ij} \right\|_\infty \lesssim n^{1/2}.$$

# Schatten classes

For simplicity,  $p = 1$ .

## Claim

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{e}_{ij} \right\|_1 \approx n^{3/2},$$

While

$$\left\| \sum_{i,j=1}^n \mathbf{e}_{ij} \right\|_1 = n.$$

The  $\geq$  side in the first equivalence follows easily from duality between  $S_1^n$  and  $S_\infty^n$  and the not-hard fact that

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{e}_{ij} \right\|_\infty \lesssim n^{1/2}.$$

# Schatten classes

Note also that for all  $\varepsilon_i, \delta_j = \pm 1$   $\|\sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{e}_{ij}\|_1 = n$ .

So, the best constant  $K$  in the inequality

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\|_1 \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|_1$$

holding for all  $\{x_{ij}\}$  in  $S_1$  is at least of order  $n^{1/2}$ .

On the other hand, it follows from Khinchine's inequality that for all  $\{x_{ij}\}$  in  $L_1$ ,

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\|_{L_1} \lesssim \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|_{L_1}. \quad (\text{upper property } \alpha)$$

It follows that the Banach–Mazur distance of  $S_1^n$  from a subspace of  $L_1$  (or any other space with “upper property  $\alpha$ ”) is at least of order  $n^{1/2}$ . It is easy to see that this is the right order.

# Schatten classes

Note also that for all  $\varepsilon_i, \delta_j = \pm 1$   $\|\sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{e}_{ij}\|_1 = n$ .  
So, the best constant  $K$  in the inequality

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\|_1 \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|_1$$

holding for all  $\{x_{ij}\}$  in  $S_1$  is at least of order  $n^{1/2}$ .

On the other hand, it follows from Khinchine's inequality that for all  $\{x_{ij}\}$  in  $L_1$ ,

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\|_{L_1} \lesssim \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|_{L_1}. \quad (\text{upper property } \alpha)$$

It follows that the Banach–Mazur distance of  $S_1^n$  from a subspace of  $L_1$  (or any other space with “upper property  $\alpha$ ”) is at least of order  $n^{1/2}$ . It is easy to see that this is the right order.

# Schatten classes

Note also that for all  $\varepsilon_i, \delta_j = \pm 1$   $\|\sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{e}_{ij}\|_1 = n$ .  
So, the best constant  $K$  in the inequality

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{x}_{ij} \right\|_1 \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{x}_{ij} \right\|_1$$

holding for all  $\{\mathbf{x}_{ij}\}$  in  $S_1$  is at least of order  $n^{1/2}$ .

On the other hand, it follows from Khinchine's inequality that for all  $\{\mathbf{x}_{ij}\}$  in  $L_1$ ,

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{x}_{ij} \right\|_{L_1} \lesssim \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{x}_{ij} \right\|_{L_1}. \quad (\text{upper property } \alpha)$$

It follows that the Banach–Mazur distance of  $S_1^n$  from a subspace of  $L_1$  (or any other space with “upper property  $\alpha$ ”) is at least of order  $n^{1/2}$ . It is easy to see that this is the right order.

Note also that for all  $\varepsilon_i, \delta_j = \pm 1$   $\|\sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{e}_{ij}\|_1 = n$ .  
So, the best constant  $K$  in the inequality

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{x}_{ij} \right\|_1 \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{x}_{ij} \right\|_1$$

holding for all  $\{\mathbf{x}_{ij}\}$  in  $S_1$  is at least of order  $n^{1/2}$ .

On the other hand, it follows from Khinchine's inequality that for all  $\{\mathbf{x}_{ij}\}$  in  $L_1$ ,

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} \mathbf{x}_{ij} \right\|_{L_1} \lesssim \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j \mathbf{x}_{ij} \right\|_{L_1}. \quad (\text{upper property } \alpha)$$

It follows that the Banach–Mazur distance of  $S_1^n$  from a subspace of  $L_1$  (or any other space with “upper property  $\alpha$ ”) is at least of order  $n^{1/2}$ . It is easy to see that this is the right order.

# non-linear embeddings

It follows from general principles (mostly differentiation) that  $c_p(S_1^n)$  is equal to their linear counterparts. But these principles no longer apply when dealing with  $c_p(A)$  for a discrete set  $A \subset S_1^n$

nor for  $c_p((S_1^n)^a)$  where for  $0 < a < 1$   $(S_1^n)^a$  denotes  $S_1^n$  with the metric  $d_a(x, y) = \|x - y\|_1^a$ .

Our purpose is to find an inequality similar to the upper property  $\alpha$  inequality but which will involve only distances between pairs of points and which holds in  $L_1$  but grossly fails in  $S_1^n$ .



It follows from general principles (mostly differentiation) that  $c_p(S_1^n)$  is equal to their linear counterparts. But these principles no longer apply when dealing with  $c_p(A)$  for a discrete set  $A \subset S_1^n$  nor for  $c_p((S_1^n)^a)$  where for  $0 < a < 1$   $(S_1^n)^a$  denotes  $S_1^n$  with the metric  $d_a(x, y) = \|x - y\|_1^a$ .

Our purpose is to find an inequality similar to the upper property  $\alpha$  inequality but which will involve only distances between pairs of points and which holds in  $L_1$  but grossly fails in  $S_1^n$ .

It follows from general principles (mostly differentiation) that  $c_p(S_1^n)$  is equal to their linear counterparts. But these principles no longer apply when dealing with  $c_p(A)$  for a discrete set  $A \subset S_1^n$  nor for  $c_p((S_1^n)^a)$  where for  $0 < a < 1$   $(S_1^n)^a$  denotes  $S_1^n$  with the metric  $d_a(x, y) = \|x - y\|_1^a$ .

Our purpose is to find an inequality similar to the upper property  $\alpha$  inequality but which will involve only distances between pairs of points and which holds in  $L_1$  but grossly fails in  $S_1^n$ .

# Digression: Enflo's type

A metric space  $(X, d_X)$  is said to have (Enflo) type  $r \in [1, \infty)$  if for every  $n \in \mathbb{N}$  and  $f : \{-1, 1\}^n \rightarrow X$ ,

$$\mathbb{E} [d_X(f(\varepsilon), f(-\varepsilon))^r] \lesssim \sum_{j=1}^n \mathbb{E} [d_X(f(\varepsilon), f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n))^r], \quad (1)$$

where the expectation is with respect to  $\varepsilon \in \{-1, 1\}^n$  chosen uniformly at random. Note that if  $X$  is a Banach space and  $f$  is the linear function given by  $f(\varepsilon) = \sum_{j=1}^n \varepsilon_j x_j$  then this is the inequality defining type  $r$ :

$$\mathbb{E} \left\| \sum_{j=1}^n \varepsilon_j x_j \right\|^r \lesssim \sum_{j=1}^n \|x_j\|^r$$

For  $p \in [1, \infty)$ ,  $L_p$  actually has Enflo type  $r = \min\{p, 2\}$ . i.e.,  $X = L_p$  satisfies (1) with  $f : \{-1, 1\}^n \rightarrow L_p$  allowed to be an arbitrary mapping rather than only a linear mapping.

# Digression: Enflo's type

A metric space  $(X, d_X)$  is said to have (Enflo) type  $r \in [1, \infty)$  if for every  $n \in \mathbb{N}$  and  $f : \{-1, 1\}^n \rightarrow X$ ,

$$\mathbb{E} [d_X(f(\varepsilon), f(-\varepsilon))^r] \lesssim \sum_{j=1}^n \mathbb{E} [d_X(f(\varepsilon), f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n))^r], \quad (1)$$

where the expectation is with respect to  $\varepsilon \in \{-1, 1\}^n$  chosen uniformly at random. Note that if  $X$  is a Banach space and  $f$  is the linear function given by  $f(\varepsilon) = \sum_{j=1}^n \varepsilon_j x_j$  then this is the inequality defining type  $r$ :

$$\mathbb{E} \left\| \sum_{j=1}^n \varepsilon_j x_j \right\|^r \lesssim \sum_{j=1}^n \|x_j\|^r$$

For  $p \in [1, \infty)$ ,  $L_p$  actually has Enflo type  $r = \min\{p, 2\}$ . i.e.,  $X = L_p$  satisfies (1) with  $f : \{-1, 1\}^n \rightarrow L_p$  allowed to be an arbitrary mapping rather than only a linear mapping.

# Digression: Enflo's type

A metric space  $(X, d_X)$  is said to have (Enflo) type  $r \in [1, \infty)$  if for every  $n \in \mathbb{N}$  and  $f : \{-1, 1\}^n \rightarrow X$ ,

$$\mathbb{E} [d_X(f(\varepsilon), f(-\varepsilon))^r] \lesssim \sum_{j=1}^n \mathbb{E} [d_X(f(\varepsilon), f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n))^r], \quad (1)$$

where the expectation is with respect to  $\varepsilon \in \{-1, 1\}^n$  chosen uniformly at random. Note that if  $X$  is a Banach space and  $f$  is the linear function given by  $f(\varepsilon) = \sum_{j=1}^n \varepsilon_j x_j$  then this is the inequality defining type  $r$ :

$$\mathbb{E} \left\| \sum_{j=1}^n \varepsilon_j x_j \right\|^r \lesssim \sum_{j=1}^n \|x_j\|^r$$

For  $p \in [1, \infty)$ ,  $L_p$  actually has Enflo type  $r = \min\{p, 2\}$ . i.e.,  $X = L_p$  satisfies (1) with  $f : \{-1, 1\}^n \rightarrow L_p$  allowed to be an arbitrary mapping rather than only a linear mapping.

# Digression: Enflo's type

A metric space  $(X, d_X)$  is said to have (Enflo) type  $r \in [1, \infty)$  if for every  $n \in \mathbb{N}$  and  $f : \{-1, 1\}^n \rightarrow X$ ,

$$\mathbb{E} [d_X(f(\varepsilon), f(-\varepsilon))^r] \lesssim \sum_{j=1}^n \mathbb{E} [d_X(f(\varepsilon), f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n))^r], \quad (1)$$

where the expectation is with respect to  $\varepsilon \in \{-1, 1\}^n$  chosen uniformly at random. Note that if  $X$  is a Banach space and  $f$  is the linear function given by  $f(\varepsilon) = \sum_{j=1}^n \varepsilon_j x_j$  then this is the inequality defining type  $r$ :

$$\mathbb{E} \left\| \sum_{j=1}^n \varepsilon_j x_j \right\|^r \lesssim \sum_{j=1}^n \|x_j\|^r$$

For  $p \in [1, \infty)$ ,  $L_p$  actually has Enflo type  $r = \min\{p, 2\}$ . i.e.,  $X = L_p$  satisfies (1) with  $f : \{-1, 1\}^n \rightarrow L_p$  allowed to be an arbitrary mapping rather than only a linear mapping.

# Digression: Enflo's type

A metric space  $(X, d_X)$  is said to have (Enflo) type  $r \in [1, \infty)$  if for every  $n \in \mathbb{N}$  and  $f : \{-1, 1\}^n \rightarrow X$ ,

$$\mathbb{E} [d_X(f(\varepsilon), f(-\varepsilon))]^r \lesssim \sum_{j=1}^n \mathbb{E} [d_X(f(\varepsilon), f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n))]^r, \quad (1)$$

where the expectation is with respect to  $\varepsilon \in \{-1, 1\}^n$  chosen uniformly at random. Note that if  $X$  is a Banach space and  $f$  is the linear function given by  $f(\varepsilon) = \sum_{j=1}^n \varepsilon_j x_j$  then this is the inequality defining type  $r$ :

$$\mathbb{E} \left\| \sum_{j=1}^n \varepsilon_j x_j \right\|^r \lesssim \sum_{j=1}^n \|x_j\|^r$$

For  $p \in [1, \infty)$ ,  $L_p$  actually has Enflo type  $r = \min\{p, 2\}$ . i.e.,  $X = L_p$  satisfies (1) with  $f : \{-1, 1\}^n \rightarrow L_p$  allowed to be an arbitrary mapping rather than only a linear mapping.

# Digression: Enflo's type

This statement was proved by Enflo in 1969 for  $p \in [1, 2]$  (and by [NS, 2002] for  $p \in (2, \infty)$ ).

Here is an illustration how to use Enflo type to show that for  $q < p \leq 2$   $c_p(\{-1, 1\}^n, \|\cdot\|_q) \gtrsim n^{\frac{1}{q} - \frac{1}{p}}$  ( $c_p(\ell_q^n) \leq n^{\frac{1}{q} - \frac{1}{p}}$  is trivial).

Let  $f : \{-1, 1\}^n \rightarrow L_p$  be such that

$$\forall x, y \in \{-1, 1\}^n, \quad \|x - y\|_q \leq \|f(x) - f(y)\|_p \leq D\|x - y\|_q$$

Then

$$2^p n^{p/q} \leq \mathbb{E} \|f(\varepsilon) - f(-\varepsilon)\|_p^p \lesssim \sum_{j=1}^n \mathbb{E} \|f(\varepsilon) - f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n)\|_p^p \lesssim D^p n 2^p.$$

So  $D \gtrsim n^{\frac{1}{q} - \frac{1}{p}}$ .



# Digression: Enflo's type

This statement was proved by Enflo in 1969 for  $p \in [1, 2]$  (and by [NS, 2002] for  $p \in (2, \infty)$ ).

Here is an illustration how to use Enflo type to show that for  $q < p \leq 2$   $c_p(\{-1, 1\}^n, \|\cdot\|_q) \gtrsim n^{\frac{1}{q} - \frac{1}{p}}$  ( $c_p(\ell_q^n) \leq n^{\frac{1}{q} - \frac{1}{p}}$  is trivial).

Let  $f : \{-1, 1\}^n \rightarrow L_p$  be such that

$$\forall x, y \in \{-1, 1\}^n, \quad \|x - y\|_q \leq \|f(x) - f(y)\|_p \leq D\|x - y\|_q$$

Then

$$2^p n^{p/q} \leq \mathbb{E} \|f(\varepsilon) - f(-\varepsilon)\|_p^p \lesssim \sum_{j=1}^n \mathbb{E} \|f(\varepsilon) - f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n)\|_p^p \lesssim D^p n 2^p.$$

So  $D \gtrsim n^{\frac{1}{q} - \frac{1}{p}}$ .

# Digression: Enflo's type

This statement was proved by Enflo in 1969 for  $p \in [1, 2]$  (and by [NS, 2002] for  $p \in (2, \infty)$ ).

Here is an illustration how to use Enflo type to show that for  $q < p \leq 2$   $c_p(\{-1, 1\}^n, \|\cdot\|_q) \gtrsim n^{\frac{1}{q}-\frac{1}{p}}$  ( $c_p(\ell_q^n) \leq n^{\frac{1}{q}-\frac{1}{p}}$  is trivial).

Let  $f : \{-1, 1\}^n \rightarrow L_p$  be such that

$$\forall x, y \in \{-1, 1\}^n, \quad \|x - y\|_q \leq \|f(x) - f(y)\|_p \leq D\|x - y\|_q$$

Then

$$2^p n^{p/q} \leq \mathbb{E} \|f(\varepsilon) - f(-\varepsilon)\|_p^p \lesssim \sum_{j=1}^n \mathbb{E} \|f(\varepsilon) - f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n)\|_p^p \lesssim D^p n 2^p.$$

So  $D \gtrsim n^{\frac{1}{q}-\frac{1}{p}}$ .

# Digression: Enflo's type

This statement was proved by Enflo in 1969 for  $p \in [1, 2]$  (and by [NS, 2002] for  $p \in (2, \infty)$ ).

Here is an illustration how to use Enflo type to show that for  $q < p \leq 2$   $c_p(\{-1, 1\}^n, \|\cdot\|_q) \gtrsim n^{\frac{1}{q}-\frac{1}{p}}$  ( $c_p(\ell_q^n) \leq n^{\frac{1}{q}-\frac{1}{p}}$  is trivial).

Let  $f : \{-1, 1\}^n \rightarrow L_p$  be such that

$$\forall x, y \in \{-1, 1\}^n, \quad \|x - y\|_q \leq \|f(x) - f(y)\|_p \leq D\|x - y\|_q$$

Then

$$2^p n^{p/q} \leq \mathbb{E} \|f(\varepsilon) - f(-\varepsilon)\|_p^p \lesssim \sum_{j=1}^n \mathbb{E} \|f(\varepsilon) - f(\varepsilon_1, \dots, \varepsilon_{j-1}, -\varepsilon_j, \varepsilon_{j+1}, \dots, \varepsilon_n)\|_p^p \lesssim D^p n 2^p.$$

So  $D \gtrsim n^{\frac{1}{q}-\frac{1}{p}}$ .

The definition of non-linear cotype is more problematic. Changing the direction of the inequality in the definition of type is no good if  $f(\{-1, 1\}^n)$  is a discrete set. A good definition was sought for a long time until the following:

A metric space  $(X, d_X)$  is said to have (Mendel-Naor) cotype  $s \in [1, \infty)$  if for every  $n \in \mathbb{N}$  there is an  $m \in \mathbb{N}$  such that for all  $f : \mathbb{Z}_{2m}^n \rightarrow X$ ,

$$\sum_{j=1}^n \frac{\mathbb{E} [d_X(f(x + me_j), f(x))^s]}{m^s} \lesssim \mathbb{E} [d_X(f(x + \varepsilon), f(x))^s],$$

where the expectation is with respect to  $(x, \varepsilon) \in \mathbb{Z}_{2m}^n \times \{-1, 0, 1\}^n$  chosen uniformly at random.

The definition of non-linear cotype is more problematic. Changing the direction of the inequality in the definition of type is no good if  $f(\{-1, 1\}^n)$  is a discrete set. A good definition was sought for a long time until the following:

A metric space  $(X, d_X)$  is said to have (Mendel-Naor) cotype  $s \in [1, \infty)$  if for every  $n \in \mathbb{N}$  there is an  $m \in \mathbb{N}$  such that for all  $f : \mathbb{Z}_{2m}^n \rightarrow X$ ,

$$\sum_{j=1}^n \frac{\mathbb{E} [d_X(f(x + me_j), f(x))^s]}{m^s} \lesssim \mathbb{E} [d_X(f(x + \varepsilon), f(x))^s],$$

where the expectation is with respect to  $(x, \varepsilon) \in \mathbb{Z}_{2m}^n \times \{-1, 0, 1\}^n$  chosen uniformly at random.

# back to non-linear version of upper property $\alpha$

We are looking for a good non-linear version of the *linear* upper property  $\alpha$  inequality:

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\| \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|.$$

We denote by  $\alpha(X)$  the best  $K$  which works for all  $x_{ij}$ -s in the normed space  $X$ .

We want to find obstructions to embedding of the grid  $M_n[m]$  of all  $n \times n$  matrices whose entries have values in  $[m] = \{-m, -(m-1), \dots, m-1, m\}$  with the  $S_1$  norm (more generally the  $S_p$  norm,  $1 \leq p < 2$ ) in a Banach space  $X$  with upper property  $\alpha$ . In particular  $L_1$  (or  $L_p$ ).

Something like the following comes to mind: For all  $f : M_n[m] \rightarrow X$ ,

$$\text{Ave}_{x,y \in M_n[m]} \|f(x) - f(y)\|^p \lesssim m^p \text{Ave}_{\substack{x \in M_n[m] \\ \varepsilon, \delta \in \{-1, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

# back to non-linear version of upper property $\alpha$

We are looking for a good non-linear version of the *linear* upper  $\alpha$  inequality:

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\| \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|.$$

We denote by  $\alpha(X)$  the best  $K$  which works for all  $x_{ij}$ -s in the normed space  $X$ .

We want to find obstructions to embedding of the grid  $M_n[m]$  of all  $n \times n$  matrices whose entries have values in  $[m] = \{-m, -(m-1), \dots, m-1, m\}$  with the  $S_1$  norm (more generally the  $S_p$  norm,  $1 \leq p < 2$ ) in a Banach space  $X$  with upper property  $\alpha$ . In particular  $L_1$  (or  $L_p$ ).

Something like the following comes to mind: For all  $f : M_n[m] \rightarrow X$ ,

$$\text{Ave}_{x,y \in M_n[m]} \|f(x) - f(y)\|^p \lesssim m^p \text{Ave}_{x \in M_n[m]} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

# back to non-linear version of upper property $\alpha$

We are looking for a good non-linear version of the *linear* upper property  $\alpha$  inequality:

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\| \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|.$$

We denote by  $\alpha(X)$  the best  $K$  which works for all  $x_{ij}$ -s in the normed space  $X$ .

We want to find obstructions to embedding of the grid  $M_n[m]$  of all  $n \times n$  matrices whose entries have values in  $[m] = \{-m, -(m-1), \dots, m-1, m\}$  with the  $S_1$  norm (more generally the  $S_p$  norm,  $1 \leq p < 2$ ) in a Banach space  $X$  with upper property  $\alpha$ . In particular  $L_1$  (or  $L_p$ ).

Something like the following comes to mind: For all  $f : M_n[m] \rightarrow X$ ,

$$\text{Ave}_{x,y \in M_n[m]} \|f(x) - f(y)\|^p \lesssim m^p \text{Ave}_{x \in M_n[m]} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$



# back to non-linear version of upper property $\alpha$

We are looking for a good non-linear version of the *linear* upper  $\alpha$  inequality:

$$\mathbb{E}_{\varepsilon_{ij}=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_{ij} x_{ij} \right\| \leq K \mathbb{E}_{\varepsilon_i, \delta_j=\pm 1} \left\| \sum_{i,j=1}^n \varepsilon_i \delta_j x_{ij} \right\|.$$

We denote by  $\alpha(X)$  the best  $K$  which works for all  $x_{ij}$ -s in the normed space  $X$ .

We want to find obstructions to embedding of the grid  $M_n[m]$  of all  $n \times n$  matrices whose entries have values in  $[m] = \{-m, -(m-1), \dots, m-1, m\}$  with the  $S_1$  norm (more generally the  $S_p$  norm,  $1 \leq p < 2$ ) in a Banach space  $X$  with upper property  $\alpha$ . In particular  $L_1$  (or  $L_p$ ).

Something like the following comes to mind: For all  $f : M_n[m] \rightarrow X$ ,

$$\text{Ave}_{x,y \in M_n[m]} \|f(x) - f(y)\|^p \lesssim m^p \text{Ave}_{\substack{x \in M_n[m] \\ \varepsilon, \delta \in \{-1, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

# back to non-linear version of upper property $\alpha$

This inequality is problematic and wrong even for  $X = \mathbb{R}$  because of the summation over different regions in the right and left sides.

There are (at least) two ways one can try to overcome this: either by wrapping  $[m]$  around, i.e. regarding summation mod  $2m + 1$ . Or by some “smoothing” of the inequality, as will be explained later.

The first method leads to elegant inequalities having to do with expansion properties of a natural graph, but unfortunately we do not see a way to use them to prove our main concern: that  $M_n[m]$  with the  $S_1^n$  distance does not nicely Lipschitz embed into  $L_1$ .

The second method leads to a solution to our problem (but as we'll see the resulting inequality is not so elegant).

# back to non-linear version of upper property $\alpha$

This inequality is problematic and wrong even for  $X = \mathbb{R}$  because of the summation over different regions in the right and left sides.

There are (at least) two ways one can try to overcome this: either by wrapping  $[m]$  around, i.e. regarding summation mod  $2m + 1$ . Or by some “smoothing” of the inequality, as will be explained later.

The first method leads to elegant inequalities having to do with expansion properties of a natural graph, but unfortunately we do not see a way to use them to prove our main concern: that  $M_n[m]$  with the  $S_1^n$  distance does not nicely Lipschitz embed into  $L_1$ .

The second method leads to a solution to our problem (but as we'll see the resulting inequality is not so elegant).

## back to non-linear version of upper property $\alpha$

This inequality is problematic and wrong even for  $X = \mathbb{R}$  because of the summation over different regions in the right and left sides.

There are (at least) two ways one can try to overcome this: either by wrapping  $[m]$  around, i.e. regarding summation mod  $2m + 1$ . Or by some “smoothing” of the inequality, as will be explained later.

The first method leads to elegant inequalities having to do with expansion properties of a natural graph, but unfortunately we do not see a way to use them to prove our main concern: that  $M_n[m]$  with the  $S_1^n$  distance does not nicely Lipschitz embed into  $L_1$ .

The second method leads to a solution to our problem (but as we'll see the resulting inequality is not so elegant).

# back to non-linear version of upper property $\alpha$

This inequality is problematic and wrong even for  $X = \mathbb{R}$  because of the summation over different regions in the right and left sides.

There are (at least) two ways one can try to overcome this: either by wrapping  $[m]$  around, i.e. regarding summation mod  $2m + 1$ . Or by some “smoothing” of the inequality, as will be explained later.

The first method leads to elegant inequalities having to do with expansion properties of a natural graph, but unfortunately we do not see a way to use them to prove our main concern: that  $M_n[m]$  with the  $S_1^n$  distance does not nicely Lipschitz embed into  $L_1$ .

The second method leads to a solution to our problem (but as we'll see the resulting inequality is not so elegant).

# Binary tensor conductance of $M_n(\mathbb{Z}_m)$

$\mathbb{Z}_m$  denotes  $\{0, 1, \dots, m-1\}$  with addition mod  $m$ .

## Theorem

Let  $m, n \in \mathbb{N}$ ,  $1 \leq p < \infty$ , with  $n^6 \lesssim_p m$  and let  $X$  be a Banach space. Let  $f : M_n(\mathbb{Z}_m) \rightarrow X$  be any function. Then

$$\mathbb{E}_{x, y \in M_n(\mathbb{Z}_m)} \|f(x) - f(y)\|^p \lesssim_p \alpha(X) m^p \mathbb{E}_{\substack{x \in M_n(\mathbb{Z}_m) \\ \varepsilon, \delta \in \{0, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

If  $X$  is  $\mathbb{R}$  (or  $L_p$ ) there is no restriction on  $m$ .

## Theorem

Let  $m, n \in \mathbb{N}$ ,  $1 \leq p \leq 2$ . Let  $f : M_n(\mathbb{Z}_m) \rightarrow \mathbb{R}$  be any function. Then

$$\mathbb{E}_{x, y \in M_n(\mathbb{Z}_m)} |f(x) - f(y)|^p \lesssim_p m^p \mathbb{E}_{\substack{x \in M_n(\mathbb{Z}_m) \\ \varepsilon, \delta \in \{0, 1\}^n}} |f(x + \varepsilon \otimes \delta) - f(x)|^p.$$

# Binary tensor conductance of $M_n(\mathbb{Z}_m)$

$\mathbb{Z}_m$  denotes  $\{0, 1, \dots, m-1\}$  with addition mod  $m$ .

## Theorem

Let  $m, n \in \mathbb{N}$ ,  $1 \leq p < \infty$ , with  $n^6 \lesssim_p m$  and let  $X$  be a Banach space. Let  $f : M_n(\mathbb{Z}_m) \rightarrow X$  be any function. Then

$$\mathbb{E}_{x, y \in M_n(\mathbb{Z}_m)} \|f(x) - f(y)\|^p \lesssim_p \alpha(X) m^p \mathbb{E}_{\substack{x \in M_n(\mathbb{Z}_m) \\ \varepsilon, \delta \in \{0, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

If  $X$  is  $\mathbb{R}$  (or  $L_p$ ) there is no restriction on  $m$ .

## Theorem

Let  $m, n \in \mathbb{N}$ ,  $1 \leq p \leq 2$ . Let  $f : M_n(\mathbb{Z}_m) \rightarrow \mathbb{R}$  be any function. Then

$$\mathbb{E}_{x, y \in M_n(\mathbb{Z}_m)} |f(x) - f(y)|^p \lesssim_p m^p \mathbb{E}_{\substack{x \in M_n(\mathbb{Z}_m) \\ \varepsilon, \delta \in \{0, 1\}^n}} |f(x + \varepsilon \otimes \delta) - f(x)|^p.$$

# Binary tensor conductance of $M_n(\mathbb{Z}_m)$

$\mathbb{Z}_m$  denotes  $\{0, 1, \dots, m-1\}$  with addition mod  $m$ .

## Theorem

Let  $m, n \in \mathbb{N}$ ,  $1 \leq p < \infty$ , with  $n^6 \lesssim_p m$  and let  $X$  be a Banach space. Let  $f : M_n(\mathbb{Z}_m) \rightarrow X$  be any function. Then

$$\mathbb{E}_{x, y \in M_n(\mathbb{Z}_m)} \|f(x) - f(y)\|^p \lesssim_p \alpha(X) m^p \mathbb{E}_{\substack{x \in M_n(\mathbb{Z}_m) \\ \varepsilon, \delta \in \{0, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

If  $X$  is  $\mathbb{R}$  (or  $L_p$ ) there is no restriction on  $m$ .

## Theorem

Let  $m, n \in \mathbb{N}$ ,  $1 \leq p \leq 2$ . Let  $f : M_n(\mathbb{Z}_m) \rightarrow \mathbb{R}$  be any function. Then

$$\mathbb{E}_{x, y \in M_n(\mathbb{Z}_m)} |f(x) - f(y)|^p \lesssim_p m^p \mathbb{E}_{\substack{x \in M_n(\mathbb{Z}_m) \\ \varepsilon, \delta \in \{0, 1\}^n}} |f(x + \varepsilon \otimes \delta) - f(x)|^p.$$



## Theorem

For every normed space  $X$  and all  $n, k$  and  $m$  satisfying  $n^6 \alpha(X) \leq k \leq C \min\{m^2/(n^6 \alpha(X)), m/n^2\}$ , there is an  $M > m$  with  $M/m \rightarrow 1$  as  $n \rightarrow \infty$  such that for all  $f : \mathbb{Z}^{n^2} \rightarrow X$ ,

$$\mathbb{E}_{\substack{x \in M_n[m] \\ \varepsilon \in M_n(\{-1,1\})}} \|f(x + 8k\varepsilon) - f(x)\|^p \\ \lesssim_p k^p \alpha^p(X) \mathbb{E}_{\substack{x \in M_n[M] \\ \varepsilon, \delta \in \{-1,1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\|^p.$$

# metric upper $\alpha$ inequality

Conversely, Assume that a Banach space  $X$  satisfy the inequality,

$$\begin{aligned} \mathbb{E}_{\substack{x \in M_n[m] \\ \varepsilon \in M_n(\{-1,1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ \leq kK \mathbb{E}_{\substack{x \in M_n[M] \\ \varepsilon, \delta \in \{-1,1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \end{aligned}$$

for all functions  $f : \mathbb{Z}^{n^2} \rightarrow X$ .

Fixing  $\{y_{ij}\} \subset X$  and applying the inequality to  $f(x) = \sum_{ij} x_{ij} y_{ij}$ , we get

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \left\| \sum_{ij} \varepsilon_{ij} y_{ij} \right\| \lesssim K \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \left\| \sum_{ij} \varepsilon_i \delta_j y_{ij} \right\|$$

which implies that  $X$  has upper property  $\alpha$  with constant  $\lesssim K$ .

# metric upper $\alpha$ inequality

Conversely, Assume that a Banach space  $X$  satisfy the inequality,

$$\begin{aligned} \mathbb{E}_{\substack{x \in M_n[m] \\ \varepsilon \in M_n(\{-1,1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ \leq kK \mathbb{E}_{\substack{x \in M_n[M] \\ \varepsilon, \delta \in \{-1,1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \end{aligned}$$

for all functions  $f : \mathbb{Z}^{n^2} \rightarrow X$ .

Fixing  $\{y_{ij}\} \subset X$  and applying the inequality to  $f(x) = \sum_{ij} x_{ij} y_{ij}$ , we get

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \left\| \sum_{ij} \varepsilon_{ij} y_{ij} \right\| \lesssim K \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \left\| \sum_{ij} \varepsilon_i \delta_j y_{ij} \right\|$$

which implies that  $X$  has upper property  $\alpha$  with constant  $\lesssim K$ .

# metric upper $\alpha$ inequality

Conversely, Assume that a Banach space  $X$  satisfy the inequality,

$$\begin{aligned} \mathbb{E}_{\substack{x \in M_n[m] \\ \varepsilon \in M_n(\{-1,1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ \leq kK \mathbb{E}_{\substack{x \in M_n[M] \\ \varepsilon, \delta \in \{-1,1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \end{aligned}$$

for all functions  $f : \mathbb{Z}^{n^2} \rightarrow X$ .

Fixing  $\{y_{ij}\} \subset X$  and applying the inequality to  $f(x) = \sum_{ij} x_{ij} y_{ij}$ , we get

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \left\| \sum_{ij} \varepsilon_{ij} y_{ij} \right\| \lesssim K \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \left\| \sum_{ij} \varepsilon_i \delta_j y_{ij} \right\|$$

which implies that  $X$  has upper property  $\alpha$  with constant  $\lesssim K$ .

# metric upper $\alpha$ inequality

## Claim

For any  $n$  and  $M$  large enough with respect to  $n$ , the distortion of embedding  $M_n(M)$  with the  $S_1$  distance into a Banach space  $X$  is, at least of order  $n^{1/2}/\alpha(X)$ .

**Proof:** If  $f : M_n[M] \rightarrow X$  is such that

$$\|x - y\|_{S_1} \leq \|f(x) - f(y)\| \leq K\|x - y\|_{S_1}$$

Then, for all  $x \in M_n[m]$  and  $\varepsilon \in M_n(\{-1, 1\})$ ,

$$\|8k\varepsilon\|_{S_1} \leq \|f(x + 8k\varepsilon) - f(x)\|_X.$$

So,

$$\begin{aligned} 8k\mathbb{E}_{\varepsilon \in M_n(\{-1, 1\})} \|\varepsilon\| &\leq \mathbb{E}_{\substack{x \in M_n[m], \\ \varepsilon \in M_n(\{-1, 1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ &\lesssim k\alpha(X) \mathbb{E}_{\substack{x \in M_n[M], \\ \varepsilon, \delta \in \{-1, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \leq kK\alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1, 1\}^n} \|\varepsilon \otimes \delta\| \end{aligned}$$

## Claim

For any  $n$  and  $M$  large enough with respect to  $n$ , the distortion of embedding  $M_n(M)$  with the  $S_1$  distance into a Banach space  $X$  is, at least of order  $n^{1/2}/\alpha(X)$ .

**Proof:** If  $f : M_n[M] \rightarrow X$  is such that

$$\|x - y\|_{S_1} \leq \|f(x) - f(y)\| \leq K\|x - y\|_{S_1}$$

Then, for all  $x \in M_n[m]$  and  $\varepsilon \in M_n(\{-1, 1\})$ ,

$$\|8k\varepsilon\|_{S_1} \leq \|f(x + 8k\varepsilon) - f(x)\|_X.$$

So,

$$\begin{aligned} 8k\mathbb{E}_{\varepsilon \in M_n(\{-1, 1\})} \|\varepsilon\| &\leq \mathbb{E}_{\substack{x \in M_n[m], \\ \varepsilon \in M_n(\{-1, 1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ &\lesssim k\alpha(X) \mathbb{E}_{\substack{x \in M_n[M], \\ \varepsilon, \delta \in \{-1, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \leq kK\alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1, 1\}^n} \|\varepsilon \otimes \delta\| \end{aligned}$$

## Claim

For any  $n$  and  $M$  large enough with respect to  $n$ , the distortion of embedding  $M_n(M)$  with the  $S_1$  distance into a Banach space  $X$  is, at least of order  $n^{1/2}/\alpha(X)$ .

**Proof:** If  $f : M_n[M] \rightarrow X$  is such that

$$\|x - y\|_{S_1} \leq \|f(x) - f(y)\| \leq K\|x - y\|_{S_1}$$

Then, for all  $x \in M_n[m]$  and  $\varepsilon \in M_n(\{-1, 1\})$ ,

$$\|8k\varepsilon\|_{S_1} \leq \|f(x + 8k\varepsilon) - f(x)\|_X.$$

So,

$$\begin{aligned} 8k\mathbb{E}_{\varepsilon \in M_n(\{-1, 1\})} \|\varepsilon\| &\leq \mathbb{E}_{\substack{x \in M_n[m], \\ \varepsilon \in M_n(\{-1, 1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ &\lesssim k\alpha(X) \mathbb{E}_{\substack{x \in M_n[M], \\ \varepsilon, \delta \in \{-1, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \leq kK\alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1, 1\}^n} \|\varepsilon \otimes \delta\| \end{aligned}$$

## Claim

For any  $n$  and  $M$  large enough with respect to  $n$ , the distortion of embedding  $M_n(M)$  with the  $S_1$  distance into a Banach space  $X$  is, at least of order  $n^{1/2}/\alpha(X)$ .

**Proof:** If  $f : M_n[M] \rightarrow X$  is such that

$$\|x - y\|_{S_1} \leq \|f(x) - f(y)\| \leq K\|x - y\|_{S_1}$$

Then, for all  $x \in M_n[m]$  and  $\varepsilon \in M_n(\{-1, 1\})$ ,

$$\|8k\varepsilon\|_{S_1} \leq \|f(x + 8k\varepsilon) - f(x)\|_X.$$

So,

$$\begin{aligned} 8k \mathbb{E}_{\varepsilon \in M_n(\{-1, 1\})} \|\varepsilon\| &\leq \mathbb{E}_{\substack{x \in M_n[m], \\ \varepsilon \in M_n(\{-1, 1\})}} \|f(x + 8k\varepsilon) - f(x)\| \\ &\lesssim k\alpha(X) \mathbb{E}_{\substack{x \in M_n[M], \\ \varepsilon, \delta \in \{-1, 1\}^n}} \|f(x + \varepsilon \otimes \delta) - f(x)\| \leq kK\alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1, 1\}^n} \|\varepsilon \otimes \delta\| \end{aligned}$$



So,

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\| \lesssim K \alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|$$

But  $\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\|_{S_1} \approx n^{3/2}$

and  $\mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|_{S_1} \approx n$ .

So  $K \gtrsim n^{1/2} / \alpha(X)$ .

So,

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\| \lesssim K \alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|$$

But  $\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\|_{S_1} \approx n^{3/2}$

and  $\mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|_{S_1} \approx n$ .

So  $K \gtrsim n^{1/2} / \alpha(X)$ .

So,

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\| \lesssim K \alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|$$

But  $\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\|_{S_1} \approx n^{3/2}$

and  $\mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|_{S_1} \approx n$ .

So  $K \gtrsim n^{1/2} / \alpha(X)$ .

So,

$$\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\| \lesssim K \alpha(X) \mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|$$

But  $\mathbb{E}_{\varepsilon \in M_n(\{-1,1\})} \|\varepsilon\|_{S_1} \approx n^{3/2}$

and  $\mathbb{E}_{\varepsilon, \delta \in \{-1,1\}^n} \|\varepsilon \otimes \delta\|_{S_1} \approx n$ .

So  $K \gtrsim n^{1/2} / \alpha(X)$ .