

# Symmetry breaking in the Hartree-Fock jellium

David Gontier

CEREMADE, Université Paris-Dauphine


Optimal transport in DFT  
Banff, January 29th 2019

joint work with Mathieu Lewin and Christian Hainzl



## Introduction: Expected phase diagram for the 3d jellium

From Jones, Ceperley, PRL 76 (1996) and Zing, Lin, Ceperley, Phys. Rev. E 66 (2002).



# Hartree-Fock jellium

= Electrons in uniform positive background, described with Hartree-Fock.

States = one-body density matrices:  $\gamma \in \mathcal{S}(L^2(\Omega, \mathbb{C}^2))$ ,  $0 \leq \gamma \leq 1$ . We write  $\gamma = \begin{pmatrix} \gamma^{\uparrow\uparrow} & \gamma^{\uparrow\downarrow} \\ \gamma^{\downarrow\uparrow} & \gamma^{\downarrow\downarrow} \end{pmatrix}$ .

Energy:

$$\mathcal{E}^{\text{HF}}(\gamma, \rho, T) = \frac{1}{2} \text{Tr}(-\Delta \gamma) + \frac{1}{2} \iint_{\Omega^2} \frac{(\rho_\gamma(\mathbf{r}) - \rho)(\rho_\gamma(\mathbf{r}') - \rho)}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}' - \frac{1}{2} \iint_{\Omega^2} \frac{\text{tr}_{\mathbb{C}^2} |\gamma(\mathbf{r}, \mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' d\mathbf{r} - T \text{Tr}(S(\gamma))$$

where  $S(t) := -t \log(t) - (1-t) \log(1-t)$  is the entropy.

Constraint:  $\text{Tr}(\gamma) = \rho|\Omega|$ .

Thermodynamic limit:  $\Omega \rightarrow \mathbb{R}^3$ , and  $\rho$  constant  $\rightarrow E^{\text{HF}}(\rho, T)$ .

**Goal:** Study the phase diagram: features of the minimisers in the  $(\rho, T)$  plane.

## Spatial symmetry breaking

If  $\gamma(\mathbf{r}, \mathbf{r}') = \gamma(\mathbf{r} - \mathbf{r}', \mathbf{0})$ , then  $\gamma$  is invariant by translation (fluid phase).

Otherwise,  $\gamma$  breaks spatial symmetry (Wigner crystallisation).

## Spin symmetry breaking

If  $\gamma^{\uparrow\uparrow} = \gamma^{\downarrow\downarrow}$  and  $\gamma^{\uparrow\downarrow} = \gamma^{\downarrow\uparrow} = 0$ , then  $\gamma$  is paramagnetic.

Otherwise, it is (partially) ferromagnetic.

# The fluid phase

Perform the minimisation only on translational-invariant states:  $\gamma(\mathbf{r}, \mathbf{r}') = \gamma(\mathbf{r} - \mathbf{r}')$ .  
 $\implies \rho_\gamma = \rho = \gamma(\mathbf{0})$  is constant  $\implies$  the **direct term** vanishes.

**Fourier operator**,  $\gamma$  is multiplication operator in Fourier by (still denoted by  $\gamma$ )

$$\gamma(\mathbf{k}) = \begin{pmatrix} \gamma^{\uparrow\uparrow}(\mathbf{k}) & \gamma^{\uparrow\downarrow}(\mathbf{k}) \\ \gamma^{\downarrow\uparrow}(\mathbf{k}) & \gamma^{\downarrow\downarrow}(\mathbf{k}) \end{pmatrix}, \quad \gamma(\mathbf{k}) = \gamma(\mathbf{k})^*, \quad 0 \leq \gamma(\mathbf{k}) \leq \mathbb{I}_2.$$

**HF energy for fluid states**

$$\frac{1}{2(2\pi)^3} \int_{\mathbb{R}^3} k^2 \text{tr}_{\mathbb{C}^2} \gamma(\mathbf{k}) d\mathbf{k} - \frac{1}{(2\pi)^5} \iint_{(\mathbb{R}^3)^2} \frac{\text{tr}_{\mathbb{C}^2} [\gamma(\mathbf{k})\gamma(\mathbf{k}')] }{|\mathbf{k} - \mathbf{k}'|^2} d\mathbf{k} d\mathbf{k}' - \frac{T}{(2\pi)^3} \int_{\mathbb{R}^3} S(\gamma(\mathbf{k})) d\mathbf{k}.$$

**Constraints**  $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \text{tr}_{\mathbb{C}^2} \gamma(\mathbf{k}) d\mathbf{k} = \rho.$

**No-spin version**  $\gamma \rightarrow g$ , that is  $g \in L^1(\mathbb{R}^3, \mathbb{R})$ ,  $0 \leq g \leq 1$  and  $(2\pi)^{-3} \int_{\mathbb{R}^3} g = \rho.$

$$\frac{1}{2(2\pi)^3} \int_{\mathbb{R}^3} k^2 g(\mathbf{k}) d\mathbf{k} - \frac{1}{(2\pi)^5} \iint_{(\mathbb{R}^3)^2} \frac{g(\mathbf{k})g(\mathbf{k}')}{|\mathbf{k} - \mathbf{k}'|^2} d\mathbf{k} d\mathbf{k}' - \frac{T}{(2\pi)^3} \int_{\mathbb{R}^3} S(g(\mathbf{k})) d\mathbf{k}.$$

## Lemma

Any minimiser among all fluid states is of the form

$$\gamma(\mathbf{k}) = U \begin{pmatrix} g^\uparrow(\mathbf{k}) & 0 \\ 0 & g^\downarrow(\mathbf{k}) \end{pmatrix} U^* \quad \text{with } U \in \text{SU}(2).$$

**Proof:**  $\text{tr}_{\mathbb{C}^2}(UD_1U^*D_2) \leq \text{tr}_{\mathbb{C}^2}(D_1D_2)$  with  $D_1, D_2$  diagonal with ordered entries.

## Corollary

$$E^{\text{HF,fluid}}(\rho, T) = \inf_{t \in [0, 1/2]} \left\{ E_{\text{nospin}}^{\text{HF,fluid}}(t\rho, T) + E_{\text{nospin}}^{\text{HF,fluid}}((1-t)\rho, T) \right\}.$$

The best  $t \in [0, \frac{1}{2}]$  is called the **polarisation**.

## Lemma (Euler-Lagrange)

Any such minimiser  $\gamma$  must satisfy the Euler-Lagrange equation

$$\gamma = \left( 1 + e^{\beta(k^2/2 - \gamma * |\cdot|^{-2} - \mu)} \right)^{-1} \quad \text{for some Lagrange multiplier } \mu \in \mathbb{R}.$$

In particular,  $g^\uparrow$  and  $g^\downarrow$  satisfy  $g^{\uparrow/\downarrow}(\mathbf{k}) = \left( 1 + e^{\beta(k^2/2 - g^{\uparrow/\downarrow} * |\cdot|^{-2} - \mu)} \right)^{-1}$  for the same  $\mu$ .

**Remark:** Spin symmetry breaking ( $g^\uparrow \neq g^\downarrow$ ) can only happen if

- the map  $\rho \mapsto \mu(\rho, T)$  is not one-to-one;
- the equation  $g \mapsto \left( 1 + e^{\beta(k^2/2 - g * |\cdot|^{-2} - \mu)} \right)^{-1}$  has at least two fixed points.

## An important example: the $T = 0$ case.

### Lemma

At  $T = 0$ , for all  $\rho > 0$ , the no-spin energy  $E_{\text{nospin}}^{\text{fluid}}$  has a unique minimiser, which is  $g := \mathbb{1}(k^2 \leq c\rho^{3/2})$ . Hence

$$E_{\text{nospin}}^{\text{fluid}}(\rho, T = 0) = C_{\text{TF}}\rho^{5/3} - C_D\rho^{4/3},$$

and

$$\mu(\rho, T = 0) = \frac{\partial}{\partial \rho} E_{\text{nospin}}^{\text{fluid}} = \frac{5}{3}C_{\text{TF}}\rho^{2/3} - \frac{4}{3}C_D\rho^{1/3} \quad (\text{not one-to-one}).$$

Including the spin, we just need to study the map

$$t \mapsto C_{\text{TF}}\rho^{5/3}(t^{5/3} + (1-t)^{5/3}) - C_D\rho^{4/3}(t^{4/3} + (1-t)^{4/3}).$$


### Theorem (G-Lewin 2018)

There is a first order phase transition at  $\rho_c = \frac{125}{24\pi^5} \left( \frac{1}{1 + 2^{1/3}} \right)^3$  ( $r_s \approx 5.45$ ):

- For  $\rho < \rho_c$ , the minimiser is unique up to global spin rotation, and it is *pure ferromagnetic* ( $g^\downarrow = 0$ );
- For  $\rho > \rho_c$ , the minimiser is unique, and is *paramagnetic*.

The energy is continuous, and has a kink at  $\rho = \rho_c$ .

## Fluid phase diagram



### Theorem (G-Lewin 2018)

For  $T \geq C\rho^{1/3}e^{-\alpha\rho^{1/6}}$ , the minimiser for the spin-fluid energy is unique and paramagnetic.



# Spatial symmetry breaking

## Theorem (Overhauser, Phys. Rev. Lett. 4, 462 (1960))

At  $T = 0$ , the fluid minimiser is **never** a HF minimiser. Actually,

$$E^{\text{HF}}(\rho, T = 0) < E^{\text{HF,fluid}}(\rho, T = 0) - C e^{-\alpha \rho^{1/6}} \quad \text{Delyon, Bernu, Baguet, Holzmann, Phys. Rev. B 92}$$

Fluid states are **unstable** with respect to the formation of Spin Density Waves (SDW).

⇒ Much more complex phase diagram.

Phase diagram at  $T = 0$  (from Baguet, Delyon, Bernu, Holzmann, Phys. Rev. B 90 (2014))




FIG. 2. Hartree-Fock phase diagram of the 3D electron gas. Energies are in Hartree per electron.  $E_M = -0.89593/r_s$  is the Madlung energy of a polarized-bcc Wigner crystal. Full lines stand for incommensurate regime ( $Q > Q_W$ ) and dashed lines for the Wigner crystal ( $Q = Q_W$ ). Thin lines stand for the polarized gas (upper curves) and thick lines for the unpolarized gas.<sup>[14]</sup>

## Theorem (G-Hainzl-Lewin 18)

- At  $T = 0$ ,

$$\left| E^{\text{HF,fluid}}(\rho, T = 0) - E^{\text{HF}}(\rho, T = 0) \right| \leq C e^{-\alpha \rho^{1/6}}.$$

- If  $\rho \gg 1$  and  $T > C e^{-\alpha \rho^{1/6}}$ ,  $E^{\text{HF}}(\rho, T)$  has a unique minimiser, which is fluid and paramagnetic. In particular,  $E^{\text{HF}}(\rho, T) = E^{\text{HF,fluid}}(\rho, T)$ .

**Idea of the proof:** Controlled the difference with the first eigenvalue of the Schrödinger-like operator


$$H(\varepsilon) := |\Delta + 1| - \frac{\varepsilon}{|\mathbf{r}|}.$$

## Lemma (G-Hainzl-Lewin 18)


The first eigenvalue  $\lambda_1(\varepsilon)$  of  $H(\varepsilon)$  satisfies

$$-C e^{-\alpha/\sqrt{\varepsilon}} \leq \lambda_1(\varepsilon) \leq -C' e^{-\alpha'/\sqrt{\varepsilon}}.$$

## Expected Phase diagram for the HF jellium



## Expected Phase diagram for the HF jellium



**Thank you for your attention!**